

Roll No.

Total Pages : 6

MCA(6/7)/DX

5520-R

COMPUTER ORIENTED NUMERICAL AND
STATISTICAL METHODS

Paper : MCA-105

Time : Three Hours]

[Maximum Marks : 80

Note : Attempt five questions in all. Question No. 1 is compulsory.
Select one question from each unit. Non-programmable
calculator can be used.

(Compulsory Question)

1. Attempt any eight of the following :

(a) If 0.333 is the approximate value of $1/3$, then percentage error is

- (i) 0.99
- (ii) 9.9
- (iii) 0.099
- (iv) 0.0099.

(b) What is the maximum number of Negative roots of the equation ?

$$f(x) = 5x^5 - 6x^3 + 4x^2 - 7 = 0$$

- (i) 3
- (ii) 2
- (iii) 5
- (iv) 4.

(c) The convergence of which of the following is of second order ?

- (i) Bisection method
- (ii) False position method
- (iii) Newton-Raphson's method
- (iv) Iterative method of form $x = \Phi(x)$.

(d) Which of the following is false ?

- (i) $\Delta = \delta E^{-1/2}$
- (ii) $\delta^2 = \Delta^2 / (1 + \Delta)$
- (iii) $E = (\delta/2 + (1 + \delta^2/4)^{1/2})^2$
- (iv) $\Delta = \mu\delta + \delta^2/2$.

(e) The number of normal equations for fitting a parabola to the given set of data using method of least squares is

- (i) 2
- (ii) 4
- (iii) 3
- (iv) 1.

(f) When 537.261 is rounded to four significant digits then the relative error is

- (i) 0.0007259
- (ii) 0.0000729
- (iii) 0.00007295
- (iv) 0.00007259.

- (g) The value of $f(2)$ of a function $y = f(x)$ for which $f(0) = 8, f(1) = 11, f(4) = 68, f(5) = 123$ is

- (i) 28
(ii) 18
(iii) 118
(iv) 11.

- (h) Statement-1: The lowest value of Chi-square is zero and highest value is infinity.

Statement-2 : In F-test, the value of F can be positive, zero or negative.

- (i) Statement-1 and Statement-2 both are false
(ii) Statement-1 is false and Statement-2 is true
(iii) Statement-1 is true and Statement-2 is false
(iv) Statement-1 and Statement-2 both are true.
- (i) In a difference table that contains an erroneous entry, the algebraic sum of the errors in any difference column is
- (i) zero
(ii) thrice the error
(iii) the error itself
(iv) twice the error. 3×8=24

UNIT-I

2. (a) (i) Define the terms 'Error', 'Relative error' and 'Inherent error'. 3
(ii) Mention various sources of Errors. 2
(iii) Round-off and truncate the following numbers correct to four significant figures : 2
63.8543 0.0063945 83615 0.090038

- (b) Find a root of

$$3x - \cos x = 1$$

by Newton-Raphson's method correct to three decimal places. 7

3. (a) Find the smallest positive root of the equation

$$x^3 - 5x + 1 = 0$$

correct to two decimal places by the False position method. 7

- (b) Write short note on any *one* of the following :

- (i) Error in number representation and computation.
(ii) Arithmetic operations with normalized floating point numbers and their consequences. 7

UNIT-II

4. (a) Explain Euler's method for finding the solution of ordinary differential equations. 7
(b) Solve the following system of simultaneous by Gauss elimination method :

$$x_1 + x_2 + x_3 = 6$$

$$3x_1 + 3x_2 + 4x_3 = 20$$

$$2x_1 + x_2 + 3x_3 = 13$$

Use complete pivoting wherever needed. 7

5. (a) Solve the following ordinary differential equation using Taylor's series method :

$$(dy/dx) = x + xy$$

Start with $x = 1, y = 0$. Find the value of y for $x = 1.2$ taking $h = 0.1$. 7

- (b) Find the value of

$$I = \int_0^1 x^2(1+x^2)^{-1} dx$$

using

- (i) Trapezoidal rule
 (ii) Simpson's one-third rule by dividing the interval $[0, 1]$ into 4 sub-intervals. 7

UNIT-III

6. (a) Derive an expression for Lagrange's interpolation formula. Apply it to find the value of $f(5.2)$ for the table of values : 7

x	4	6	8	10
$f(x)$	19	40	79	142

- (b) For the data given below, find the equation of the best fitting exponential curve of the form $y = ae^{bx}$. 7

x	1	2	3	4	5	6
y	1.6	4.5	13.8	40.2	125	300

7. (a) What observations you make of the effect of an error in an entry in a difference table. 7
 (b) Given data below, find the equation of the best fit curve of the form $y = ax^2 + bx + c$. 7

x	1	2	3	4	5	6
y	1.6	4.5	13.8	40.2	125	300